

this case, however, the interest of the record lies in the fact that these specimens had been bred here and were taken from the larval gallery seven inches from the outside of the tree. This was testified by a piece of the wood which Mr. Dixon-Nuttall was good enough to send me displaying the galleries. A credible explanation of the origin of the progenitors of these specimens is afforded by the fact that some years ago new gate posts were put down in or near this farm, some of which were made of American ash.—W. E. SHARP, South Norwood: *March 10th*, 1905.

[The late Mr. P. B. Mason has recorded the capture of *N. erythrocephalus*, with another North American species, *N. caprea*, Say, at Burton-on-Trent, in an ash tree which had been brought from Carrick-on-Suir, Ireland. cf. Ent. Mo. Mag., vol. xxxiii, p. 91 (1897).—J. J. W].

*Anisotoma furva*, Er., at Skegness.—On Sept. 11th, 1904, at Skegness, Lincs., by searching in the hollows on the sand-hills between 5 and 6 p.m. I took amongst a host of common beetles, 5 *Anisotomas* made up of one *A. dubia*, Kugel. (small var.), one *A. ovalis*, Schm., and three of the rare *A. furva*, Er. (two ♀, one ♂). They have been examined and the names kindly supplied by Mr. G. C. Champion.—E. W. MORSE, 9, Hill Top Mount, Roundhay Road, Leeds: *March 15th*, 1905.

*Ptinus tectus*, Boield.: *Synonymic note*.—This species was first introduced by Boieldieu in his "Monographie des Ptiniore" (Ann. Ent. Soc. Fr., 1854, 652). He gives *P. pilosus*, White (Voy. Ereb. Terr., 1846, xi, 8), as a synonym, and this synonymy has been reproduced in various catalogues. The type of White's *pilosus* (which is labelled "*pilosulus*") is in the British Museum. Boieldieu's type is in the possession of M. Bedel, and he has been kind enough to carefully compare British specimens received from me with this type. Upon comparing these with White's *pilosus*, it is evident that there is not the slightest resemblance between them. White's insect is an elongate, parallel-sided insect, with close, decumbent, somewhat greenish-grey pubescence, and is from New Zealand. It is remarkable that in some points Boieldieu's description agrees better with White's insect than with my examples referred to above; indeed, the description is a bad one for what we now call *Ptinus tectus*, nor does it altogether accord with White's *pilosulus*.—E. A. NEWBERY, 12, Churchill Road, Dartmouth Park, N.W.: *February 15th*, 1905.

*Diptera in the New Forest*.—Mr. A. E. Gibbs, of St. Albans, has lately sent me for determination a number of *Diptera* collected for him by Mr. W. Brameld, of Brockenhurst. Among them are several species which may be worth mention as usually rare, though some of them are not uncommon in the New Forest. I cannot give the dates and localities, but all were taken in that district and almost all in 1904.

Of the *Nematocera* I would only mention *Limnobia annulus*, Mg. A fine and very local species, and *Pedicia rivosa*, L., not perhaps uncommon, but a large and very handsome insect.

Of the *Brachycera*—*Atylotus fulvus*, Mg., and *Chrysops quadrata*, Mg., seem to have been common in the Forest, as well as several other *Tabanidæ*. To these may be added the pretty *Oxycera pulchella*, Mg, the exotic looking *Anthrax fenest-*

*tratus*, Fln., and the strange little *Oncodes gibbosus*, L. The *Syrphidæ* were well represented, though mostly by common species; the following, however, are usually scarce, *Melangyna quadrimaculata*, Verr., *Xanthogramma citrofasciatum*, Deg., of this there was only one specimen, though there were several of the much commoner *X. ornatum*, Mg., *Eristalis cryptarum*, F., ♂ and ♀, *Xylota lenta*, Mg., and *Chrysotoxum elegans*, Lw.; this latter though usually scarce would seem to be not uncommon in the Forest.

The best *Conopidæ* were *Conops vesicularis*, L., ♂ and ♀, and *C. ceriiformis* Mg., the former used to be considered a great rarity, but is apparently not so now. To these I would add *Hypoderma? lineatum*, Vill., probably common in the larva state, the two fine Tachinids, *Echinomyia grossa*, L., and *Alophora hemiptera*, F., neither of which seem scarce, *Helomyza pectoralis*, Lw., *Phæomyia fuscipennis*, Mg., *Pteropæctria afflicta*, Mg., and *P. palustris*, Mg., and last, but not least interesting, the rare *Icterica westermanni*, Mg., of which ten specimens were taken by sweeping rushes at Milford soon after harvest.

Mr. Andrews has kindly sent me a list of species from the New Forest, taken during the last and previous seasons. The following seem worthy of mention, besides those given in his note (*Ent. Mo. Mag.*, March, 1905, p. 71), *Xanthandrus comtus*, Harr., *Didea fasciata*, Meq., *D. intermedia*, Lw., *Tolucella inanis*, L. (taken by Mr. Brameld), *Mallota cimbiciformis*, Fln., and *Mgiolepta luteola*, Gmel.—E. N. BLOOMFIELD, Guestling: *March*, 1905.

*Rhanphomyia tenuirostris*, Fal., taken in the New Forest.—Among some *Diptera* recently received from Mr. Cartor was a specimen of this species, taken in Arran, and on placing it in my cabinet I was surprised to find three specimens under this name, about which I had quite forgotten. On referring to my catalogue they proved to have been taken at Lyndhurst; two in September, 1900, and the other in September, 1901; and there is also a note showing that at the time I was doubtful as to the name being correct, as this species is not included in the British list. Mr. Grimshaw, in his "*Diptera Scotica*," records a single female taken at Glencorse, September 8th, 1898, and adds: "The only other British record of this species with which I am acquainted is that given by Col. Yerbury in the '*Irish Naturalist*,' March, 1902," where he mentions a specimen taken at Loo Bridge, in Ireland." Curtis, however, recorded it about the year 1825 from a female specimen taken in the Isle of Wight, so although it may not be often met with, it is evidently widely distributed. I am informed Mr. Grimshaw considers the generic name, *Macrostomus*, Wied., has priority.—F. C. ADAMS, 50, Ashley Gardens, S.W.: *March 2nd*, 1905.

*Dr. Reuter on the Urostylinæ*.—In his interesting remarks on this subfamily of the *Pentatomidæ* (*ante*, p. 64), Dr. Reuter—who follows Dallas in considering that the *Urostylinæ* constitute a distinct family—has made some reference to my first volume on the *Rhynchota* of British India, which may perhaps create a wrong impression. His remarks may be taken to suggest that I have not noticed his genus *Eurhynchocoris*, which he described in 1881. This, however, is not the case; at the foot of my *Synopsis* of the genera I have added the following note:—"The